

# FRIED'S THEOREM FOR BOUNDARY GEOMETRIES OF RANK ONE SYMMETRIC SPACES

Raphaël V. ALEXANDRE\*

September 22, 2019

## Abstract

After introducing the different boundary geometries of rank one symmetric spaces, we state and prove Fried's theorem in the general setting of all those geometries: a closed manifold with a similarity structure is either complete or the developing map is a covering onto the Heisenberg-type space deprived of a point.

## I — INTRODUCTION

Let  $\mathbf{F}$  be the field of the real, complex, quaternionic or octonionic numbers. We are interested in the boundary geometries  $(\mathrm{PU}(n, 1; \mathbf{F}), \partial\mathbf{H}_{\mathbf{F}}^n)$ . Those structures will be considered for  $n \geq 2$ . For the non-real case, the hyperbolic lines (when  $n = 1$ ) are isometric to real hyperbolic spaces, so the hypothesis  $n \geq 2$  is only a convenience. When  $\mathbf{F}$  is the octonionic field, we only consider the case  $n = 2$ . For example, when  $\mathbf{F} = \mathbf{R}$ , we get the flat conformal structure. When  $\mathbf{F} = \mathbf{C}$ , we get the spherical CR structure.

In this paper we will prove the following theorem, called Fried's theorem since it was stated and proved by Fried for the real case in [Fri80]. It was also proved for the complex case by Miner [Min90] and for the quaternionic case by Kamishima [Kam99]. It seems that the octonionic case has not been proved yet.

Start with a rank one symmetric space. We set  $\mathcal{N}$  the geodesic boundary of the space deprived of a point and  $\mathrm{Sim}(\mathcal{N})$  the subgroup of the isometries stabilizing this point.

**THEOREM (III.1)** *Let  $M$  be a closed  $(\mathrm{Sim}(\mathcal{N}), \mathcal{N})$ -manifold. If the developing map  $D: \tilde{M} \rightarrow \mathcal{N}$  is not a cover onto  $\mathcal{N}$ , then the holonomy subgroup  $\Gamma$  fixes a point in  $\mathcal{N}$  and  $D$  is in fact a covering onto the complement of this point.*

The proof proposed here of Fried's theorem III.1 will simultaneously deal with all the cases. This unified proof will use a general approach of convexity in  $\mathcal{N}$ . Convexity arguments were crucial in Fried's initial approach. In the real case, the space and the tangent space on one hand, and the geodesic structure and the algebraic sum on the other hand, are each time essentially the same. Therefore a clarification was required to state general convexity arguments for every field.

A consequence of Fried's theorem is the following result. We keep the notations of the preceding theorem and denote by  $L(\Gamma)$  the limit set of the holonomy group  $\Gamma$ .

**THEOREM (IV.1)** *Let  $M$  be a closed  $(\mathrm{PU}(n, 1; \mathbf{F}), \partial\mathbf{H}_{\mathbf{F}}^n)$ -manifold. If  $D$  is not surjective then it is a covering onto its image. Furthermore,  $D$  is a covering on its image if, and only if,  $D(\tilde{M})$  is equal to a connected component of  $\partial\mathbf{H}_{\mathbf{F}}^n - L(\Gamma)$ .*

---

\*Institut de Mathématiques de Jussieu-Paris Rive Gauche, Sorbonne Université, 4 Place Jussieu, 75252 Paris Cédex, France. Email address: raphael.alexandre@imj-prg.fr

The first section being the present introduction, the second section will introduce the different notions required. Notations for  $(G, X)$ -structures will be introduced in II.5. The third section consists of the proof of Fried's theorem. The forth consists of the proof of theorem IV.1.

## CONTENTS

<b>I</b>	<b>Introduction</b>	<b>1</b>
<b>II</b>	<b>Hyperbolic geometry</b>	<b>2</b>
II.1	The $KAN$ -Iwasawa decomposition	3
II.2	The group $N$	4
II.3	The octonionic case	5
II.4	Limit sets	5
II.5	Complete structures	6
<b>III</b>	<b>Fried's theorem and similarity structures</b>	<b>7</b>
III.1	A shortcut: discrete holonomy and autosimilarity	7
III.2	The geometry of $\mathcal{N}$	8
III.3	Proof of Fried's theorem	11
<b>IV</b>	<b>Complete structures on closed manifolds</b>	<b>18</b>

## II — HYPERBOLIC GEOMETRY

For the moment we will set the octonionic case aside. Let  $\mathbf{F}$  be the field of the real or complex or quaternionic numbers. We denote by  $\mathbf{F}^{n,1}$  the space  $\mathbf{F}^{n+1}$  endowed with the quadratic form

$$Q_E(z_1, \dots, z_{n+1}) = z_1 z_1^* + \dots + z_n z_n^* - z_{n+1} z_{n+1}^*, \quad (1)$$

where  $z^*$  denotes the conjugated of  $z$ . We specify  $E$  in  $Q_E$  to indicate that  $Q$  is given in the canonical basis  $(e_1, \dots, e_{n+1})$  of  $\mathbf{F}^{n+1}$ .

Now, let  $F$  be the basis

$$f_1 = \frac{-e_1 + e_{n+1}}{\sqrt{2}}, f_2 = e_2, \dots, f_n = e_n, f_{n+1} = \frac{e_1 + e_{n+1}}{\sqrt{2}}. \quad (2)$$

It is clear that it is an orthonormal basis with respect to  $Q$ . In that basis,  $Q_F$  is given by

$$Q_F(w_1, \dots, w_{n+1}) = w_2 w_2^* + \dots + w_n w_n^* - 2\operatorname{Re}(w_1 w_{n+1}^*). \quad (3)$$

Here, we denote by  $\operatorname{Re}(p)$  and  $\operatorname{Im}(p)$  the real and imaginary parts of  $p$  such that  $p = \operatorname{Re}(p) + \operatorname{Im}(p)$  and  $\operatorname{Re}(p) \in \mathbf{R}$  and  $\operatorname{Im}(p) \in \operatorname{Im}(\mathbf{F}) = \operatorname{Re}^{-1}(0)$ .

We denote by  $U(k)$  the group of the unitary matrices of  $\mathbf{F}^k$ . We denote by  $U(n, 1)$  the group of the unitary matrices of  $\mathbf{F}^{n,1}$ , i.e. the matrices acting by isometries on  $\mathbf{F}^{n,1}$ . We denote by  $\mathbf{H}^n$  the projectivized of the negative subspace  $\{Q(p) < 0\}$  and by  $\partial\mathbf{H}^n$  the projectivized null subspace  $\{Q(p) = 0\}$ . Of course topologically speaking,  $\partial\mathbf{H}^n$  is in fact the boundary of  $\mathbf{H}^n$ . The image of  $U(n, 1)$  in  $\operatorname{PGL}(\mathbf{F}^{n+1})$  is denoted by  $\operatorname{PU}(n, 1)$ .

We recover the ball-model of  $\mathbf{H}^n$  as follows. In  $\mathbf{P}(\mathbf{F}^{n+1})$  we take the affine chart  $z_{n+1} = 1$ . Thus,

$$Q_E(z) \leq 0 \iff |z_1|^2 + \dots + |z_n|^2 \leq 1. \quad (4)$$

In what follows we will take these coordinates of  $\mathbf{H}^n$ . In  $\mathbf{H}^n$ , the vector  $f_1$  represents “ $-1$ ” =  $[-1, 0, \dots, 0, 1]$  and  $f_{n+1}$  represents “ $1$ ” =  $[1, 0, \dots, 0, 1]$  which are in  $\partial\mathbf{H}^n$ . In the basis  $E$ , the vector  $e_{n+1}$  represents “ $0$ ” =  $[0, \dots, 0, 1]$  and is in  $\mathbf{H}^n$ . The other vectors of the basis  $E$  and  $F$  are not in  $\mathbf{H}^n$  nor in  $\partial\mathbf{H}^n$ .

## II.1 THE $KAN$ -IWASAWA DECOMPOSITION

We will now always suppose that  $n \geq 2$ .

We denote by  $K$  the subgroup of  $\mathrm{PU}(n, 1)$  given by the matrices in the canonical basis

$$k = \begin{pmatrix} k' & \\ & 1 \end{pmatrix} \quad (5)$$

where  $k'$  belongs to  $\mathrm{U}(n)$ .

We denote by  $\mathrm{U}(\mathcal{N})$  or also  $M$  the subgroup of  $K$  stabilizing  $\mathbf{F}f_1$  and  $\mathbf{F}f_{n+1}$ . In general, see [KP03], the matrices of  $M$  are given in the basis  $F$  by

$$m = \begin{pmatrix} \alpha & & \\ & \alpha m' & \\ & & \alpha \end{pmatrix} \quad (6)$$

with  $m' \in \mathrm{U}(n-1)$  and  $\alpha \in \mathrm{U}(1)$ . Since we projectivize on the right, the unitary factor  $\alpha$  can only be forgotten in the cases  $\mathbf{F} = \mathbf{R}$  or  $\mathbf{F} = \mathbf{C}$ .

We denote by  $A$  the subgroup of  $\mathrm{PU}(n, 1)$  given by the matrices in the basis  $F$

$$a_t = \begin{pmatrix} e^{-t} & & \\ & E_{n-1} & \\ & & e^t \end{pmatrix} \quad (7)$$

where  $t \in \mathbf{R}$  and  $E_{n-1}$  is the identity matrix. We denote by  $N$  the subgroup of  $\mathrm{PU}(n, 1)$  given by the matrices in the basis  $F$

$$n_{u,I} = \begin{pmatrix} 1 & & & \\ u & E_{n-1} & & \\ \|u\|^2/2 + I & {}^t u^* & & 1 \end{pmatrix} \quad (8)$$

where  $(u, I) \in \mathbf{F}^{n-1} \times \mathrm{Im}(\mathbf{F})$ . The vector  $u$  is to be thought as in  $(1, u, 1)$  in  $\mathbf{F}^{n,1}$ .

We denote by  $P$  the product  $MAN$ . Classical linear algebra shows the following lemma.

**LEMMA II.1** *The following properties are true.*

- (1) *The action of  $N$  is free and transitive on  $\partial\mathbf{H}^n - \{1\}$ .*
- (2) *The subgroup  $K$  is the stabilizer of  $0$  and is transitive on  $\partial\mathbf{H}^n$ .*
- (3) *We have  $\mathrm{PU}(n, 1) = KP = KAN$ .*
- (4) *The subgroup  $P$  is the stabilizer of  $1$  and is transitive on  $\mathbf{H}^n$ .*

## II.2 THE GROUP $N$

We will now closely study the subgroup  $N \subset \text{PU}(n, 1)$ . First, we identify  $n_{u,I}$  with the couple  $(u, I) \in \mathbf{F}^{n-1} \times \text{Im}(\mathbf{F})$ . We have:

$$\begin{aligned}
 (u, I) + (v, J) &:= n_{u,I} n_{v,J} \\
 &= \begin{pmatrix} 1 & & & \\ & u+v & & \\ & \|u\|^2/2 + I + u^*v + \|v\|^2/2 + J & E_{n-1} & \\ & {}^t(u+v)^* & & 1 \end{pmatrix} \\
 &= \begin{pmatrix} 1 & & & \\ & u+v & & \\ & \|u+v\|^2/2 + \text{Im}(u^*v) + I + J & E_{n-1} & \\ & {}^t(u+v)^* & & 1 \end{pmatrix} \\
 &= (u+v, I+J+\text{Im}(u^*v)).
 \end{aligned} \tag{9}$$

Indeed, computations show that

$$\text{Im}(u)\text{Im}(v) = \text{Re}(u)\text{Re}(v) + \text{Re}(u)\text{Im}(v) - \text{Re}(v)\text{Im}(u) - u^*v \tag{10}$$

$$\text{Im}(u)\text{Im}(v) + \text{Im}(v)\text{Im}(u) = 2\text{Re}(u)\text{Re}(v) - 2\text{Im}(u^*v) \tag{11}$$

$$\begin{aligned}
 \|u+v\|^2 &= \text{Re}(u)^2 + 2\text{Re}(u)\text{Re}(v) - \text{Im}(u)^2 - \text{Im}(u)\text{Im}(v) \\
 &\quad + \text{Re}(v)^2 - \text{Im}(v)\text{Im}(u) - \text{Im}(v)^2
 \end{aligned} \tag{12}$$

$$= \|u\|^2 + \|v\|^2 + 2\text{Re}(u^*v) \tag{13}$$

$$\|u\|^2/2 + \|v\|^2/2 + u^*v = \|u+v\|^2/2 + \text{Im}(u^*v). \tag{14}$$

This addition is not commutative, since  $\text{Im}(u^*v)$  is not symmetric. We have  $-(u, I) = (-u, -I)$ . We also have additivity since

$$\text{Im}(u^*v) + \text{Im}((u+v)^*w) = \text{Im}(u^*v) + \text{Im}(u^*w) + \text{Im}(v^*w) \tag{15}$$

$$= \text{Im}(u^*(v+w)) + \text{Im}(v^*w). \tag{16}$$

By the action of  $P$  on  $\partial\mathbf{H}^n$ , the point 1 is fixed. Also, the action of  $N$  on the point  $-1$  is free and transitive on  $\partial\mathbf{H}^n - \{1\}$  (see lemma II.1). We denote by  $\mathcal{N}$  the image of  $-1$  by the correspondance  $n_{u,s} \leftrightarrow (u, s)$ . Thus, we geometrically obtain  $\mathcal{N} = \partial\mathbf{H}^n - \{1\}$  and also  $\mathcal{N}$  isomorphic to  $N$ . Note that,  $-1 \in \partial\mathbf{H}^n$  corresponds to  $0 \in \mathcal{N}$  and  $1 \in \partial\mathbf{H}^n$  to  $\infty \in \mathcal{N}$ .

Now, we look at the action of  $M$  and  $A$  on  $N$  by looking of the action on  $-1$ . Easy calculations show that

$$mn_{u,I}(-1) = n_{\alpha m' u \alpha^{-1}, \alpha I \alpha^{-1}}(-1), \tag{17}$$

$$a_t n_{u,I}(-1) = n_{e^t u, e^{2t} I}(-1). \tag{18}$$

Thus, we can see that the correspondance  $n_{u,I} \leftrightarrow (u, I)$  gives  $a_t n_{u,I} \leftrightarrow (e^t u, e^{2t} I)$  and  $mn_{u,I} \leftrightarrow (\alpha m' u \alpha^{-1}, \alpha I \alpha^{-1})$ . Changing  $e^t$  for  $\lambda$  and  $(\alpha m', \alpha)$  for  $(P, \alpha)$ , denoting  $\lambda := a_t$  and  $P := m$ , we get the actions:

$$P(u, I) = (Pu\alpha^{-1}, \alpha I \alpha^{-1}) \tag{19}$$

$$\lambda(u, I) = (\lambda u, \lambda^2 I). \tag{20}$$

Remark that the actions of  $A$  and  $M$  commute. If  $f \in \text{Sim}(\mathcal{N})$ , then we can express  $f$  as

$$f(x) = \lambda P(x) + c \tag{21}$$

with  $\lambda \in \mathbf{R}_+$ ,  $P \in \text{U}(\mathcal{N})$  and  $c \in \mathcal{N}$ .

Now we give a pseudo-norm on  $\mathcal{N}$  (compare with [Cow+91, p. 10]). For  $x = (u, I) \in \mathcal{N}$  we define

$$\|x\|_{\mathcal{N}}^2 := \|u\|_{\mathbf{F}^{n-2}}^2 + \|I\|_{\mathbf{F}}^2 \quad (22)$$

It is easy to see that  $\|x\| = 0$  if and only if  $x = (0, 0)$  and that

$$\|\lambda x\|_{\mathcal{N}} = \lambda \|x\|_{\mathcal{N}} \quad (23)$$

for any  $\lambda \in \mathbf{R}_+$ . The triangle inequality is respected.

To  $\|\cdot\|_{\mathcal{N}}$  we can associate the distance function

$$d_{\mathcal{N}}(x, y) = \|-x + y\| = d_{\mathcal{N}}(0, -x + y). \quad (24)$$

and remark that it is symmetric since  $\|-(u, I)\| = \|(-u, -I)\| = \|(u, I)\|$  and  $\|-y + x\| = \|-(-x + y)\|$ .

### II.3 THE OCTONIONIC CASE

The special case  $\mathbf{F} = \mathbf{O}$  needs extra care (see [Bae02] for a general survey on the octonions). Indeed,  $\mathbf{O}^3$  is not a vector space and the preceding construction relies on this structure. However, as explained by Allcock in [All99] (compare also with [Har90] for the projective plane), most of the construction can still be made.

We can still speak of the hyperbolic plane  $\mathbf{H}_{\mathbf{O}}^2$  as a smooth manifold. It can even be made from elements of  $\mathbf{O}^3$  such that the coordinates lie in an associative algebra [All99, p. 12] (we have different conventions).

Again,  $\text{PU}(2, 1; \mathbf{O}) := \text{Aut}(\mathbf{H}_{\mathbf{O}}^2)$  acts transitively on the boundary at infinity (which can still be defined)  $\partial\mathbf{H}_{\mathbf{O}}^2$ . The 15-dimensional subgroup  $\mathbf{N}$  stabilizing a point  $\infty \in \partial\mathbf{H}_{\mathbf{O}}^2$  is similar to the Heisenberg group (it is a  $H$ -type group – see [Cow+91] for an algebraic approach), hence giving the space  $\mathcal{N}$  by the same identification. Translations are again given by

$$(x, z) + (\xi, \eta) = (x + \xi, z + \eta + \text{Im}(x^* \eta)). \quad (25)$$

Here, the coordinates  $(x, z)$  are as previously with  $x \in \mathbf{O}$  and  $z \in \text{Im}(\mathbf{O})$ . For the same reason as for the other fields, the addition of  $\mathcal{N}$  remains associative since the addition in  $\mathbf{O}$  is associative, and the addition in  $\mathcal{N}$  is not commutative. As in the general case, the commutator of two translations is given by  $(0, 2\text{Im}(x^* \eta))$ . If  $\mu$  is a unit imaginary, then we can define a rotation by  $\mu$  on  $\mathcal{N}$ .

$$m_{\mu}(x, z) = (\mu x, \mu z \mu^{-1}). \quad (26)$$

It gives a compact group of rotations. If  $\lambda \in \mathbf{R}_+$ , then again we have dilatations given by

$$\lambda(x, z) = (\lambda x, \lambda^2 z). \quad (27)$$

So the group  $\text{Sim}(\mathcal{N})$  is very similar to the previous cases. We also remark that the stabilizer  $K$  of a point in the interior  $0 \in \mathbf{H}_{\mathbf{O}}^2$  is still a compact group and again  $\mathbf{H}_{\mathbf{O}}^2 = \text{Aut}(\mathbf{H}_{\mathbf{O}}^2)/K$ . Finally, the same norm  $\|\cdot\|_{\mathcal{N}}$  can be defined and used as previously.

More references can be given. In [Rie82], Riehm shows that the isometry group of  $\partial\mathbf{H}_{\mathbf{O}}^2$  stabilizing  $\infty$  is transitive on the unit sphere. In [Rie84], Riehm shows that every geodesic is a one-parameter group of global isometries. Those results and a full description of the automorphism group are described in [Cow+91]. Note that those results are common for every field  $\mathbf{F}$  we consider here.

### II.4 LIMIT SETS

In what follows we will use limit sets (compare with [CG74]). Given a subgroup  $\Gamma \subset \text{PU}(n, 1)$ , we can define the *limit set of*  $\Gamma$ ,  $L(\Gamma)$ , as  $\overline{\Gamma \cdot p} \cap \partial\mathbf{H}^n$  for any  $p \in \mathbf{H}^n$ . This

does not depend on the choice of  $p$  since for  $q \in \mathbf{H}^n$ , the distance between  $p$  and  $q$  is preserved by elements of  $\Gamma$ . Hence, if  $g_n p \rightarrow x \in \partial \mathbf{H}^n$ , then so must  $g_n q$ .

Also, for  $x \in L(\Gamma)$  such as  $x = \lim g_n p$ , any other point  $z$  of  $\partial \mathbf{H}^n$  except possibly one must verify  $\lim g_n z = x$ . Indeed, given  $z_1, z_2$  different from one another and both distinct from  $x$ , then for  $\gamma$  a geodesic between  $z_1, z_2$ , since the points of the geodesic go to  $x$ , then so must one extremity of the geodesic. If  $y$  is the only point that does not tend to  $x$ , then we call  $(x, y)$  a *dual pair*, or *dual points*. In this case, it is easy to see that  $\lim g_n^{-1} p = y$ .

The limit set  $L(\Gamma)$  is stable by the action of  $\Gamma$ . Moreover, it is the minimal invariant set by  $\Gamma$ : if  $A \subset \partial \mathbf{H}^n$  is *closed and invariant by  $\Gamma$  and is at least constituted of two points*, then  $L(\Gamma) \subset A$ . This fact can be deduced from the preceding remark.

The following lemma will be used in what follows. The proof is easily extended to the case where  $\mathbf{F}$  is the octonion field. It uses the fact that  $K$  in the decomposition  $KAN$  is maximal compact. Another way to deduce this lemma is given by CAT(0) theory (see [BH99, p. 179]).

**LEMMA II.2** (See [CG74, p. 79]) *Let  $\Gamma \subset \mathrm{PU}(n, 1)$  be a subgroup. Suppose  $L(\Gamma) = \emptyset$ , then the elements of  $\Gamma$  let a point fixed in  $\mathbf{H}^n$ .*

## II.5 COMPLETE STRUCTURES

A  $(G, X)$ -structure (compare with Thurston's textbook [Thu97]), is a pair of a smooth space  $X$  with a transitive group  $G$  acting by analytic diffeomorphisms. (A more general notion in [Thu97] appears by allowing diffeomorphisms to be only locally defined, but we won't need this generality here.) A smooth manifold  $M$  gets a  $(G, X)$ -structure if we can choose an atlas of  $M$  consisting of charts defined on open sets of  $X$  and with transition maps belonging to  $G$ . In this case we speak of a  $(G, X)$ -manifold to designate a manifold together with a  $(G, X)$ -structure.

Such a  $(G, X)$ -structure on a smooth manifold  $M$  gives a pair  $(D, \rho)$  of the *developing map*  $D: \tilde{M} \rightarrow X$  and the *holonomy map*  $\rho: \pi_1(M) \rightarrow G$ . The developing map is a local diffeomorphism. The image of the holonomy map is called the *holonomy group*, and is usually denoted here by  $\Gamma = \rho(\pi_1(M))$ . This pair  $(D, \rho)$  prescribes the  $(G, X)$ -structure on  $M$ .

Two general problems naturally arise: it is hard to say when it is possible to put a  $(G, X)$ -structure on  $M$  (it is the *geometrization* problem); and there are very few general properties on a pair  $(D, \rho)$ . Two properties on  $(D, \rho)$  are interesting to investigate: the completeness ( $D$  is a covering), and the discreteness ( $\Gamma$  is discrete in  $G$ ).

We will say that the structure is *complete* if the developing map  $D: \tilde{M} \rightarrow X$  is a covering onto  $X$ . Of course, complete  $(\mathrm{PU}(n, 1), \partial \mathbf{H}^n)$ -manifolds are rare (those have a spherical structure), and we will rather ask if  $D: \tilde{M} \rightarrow X$  is a covering onto its image (but we keep the term *complete* for a covering onto  $X$ ). Remark that if  $D$  is a covering map onto a simply connected space, then  $\Gamma$  is discrete. (The converse is not true in general, see for example [Fal08].)

The complete structures that we will encounter come from the following lemma. None of these results are new. (Compare again with [Thu97].)

**LEMMA II.3** *Let  $(G, X)$  be a geometrical structure such that  $X$  has a  $G$ -invariant riemannian metric. If  $M$  is a closed  $(G, X)$ -manifold, then it is complete.*

**PROOF** Let  $D: \tilde{M} \rightarrow X$  be the developing map. It is a local diffeomorphism. Pulling-back the metric on  $\tilde{M}$ , it makes  $D$  a local isometry. Since  $M$  is closed,  $D$  is a covering map. ⌘

Therefore, to show that a  $(G, X)$ -structure is complete for a closed manifold  $M$ , it suffices to show that  $G$  has compact stabilizers on  $X$  (compare with [Thu97, p. 144]).

Recall that  $K \subset \mathrm{PU}(n, 1)$  from the Iwasawa decomposition is compact. And remark that  $(\mathrm{Sim}(\mathcal{N})_0, \mathcal{N} - \{0\})$  has compact stabilizers.

**LEMMA II.4** *Any closed  $(K, \partial\mathbf{H}^n)$  or  $(\mathrm{Sim}(\mathcal{N})_0, \mathcal{N} - \{0\})$ -manifold is complete.  $\mathbb{Z}$*

**LEMMA II.5** *Let  $M$  be a closed  $(\mathrm{PU}(n, 1), \partial\mathbf{H}^n)$ -manifold. Let  $\Gamma$  be the holonomy group. If  $L(\Gamma) = \emptyset$ , then  $D: \tilde{M} \rightarrow \partial\mathbf{H}^n$  is a covering map (therefore a diffeomorphism).*

**PROOF** If  $L(\Gamma) = \emptyset$ , then the elements of  $\Gamma$  let a point fixed in  $\mathbf{H}^n$  by lemma II.2. Up to conjugation it is 0, hence  $\Gamma \subset K$ . By lemma II.4, it follows that  $D$  is a covering map.  $\mathbb{Z}$

**COROLLARY II.6** *Let  $M$  be a closed  $(\mathrm{PU}(n, 1), \partial\mathbf{H}^n)$ -manifold. If  $L(\Gamma) = \emptyset$ , then  $M$  is a spherical manifold. In particular, if  $M$  is a simply connected closed  $(\mathrm{PU}(n, 1), \partial\mathbf{H}^n)$ -manifold, then  $M$  is diffeomorphic to a sphere.*

**PROOF** Because  $K \simeq \mathrm{U}(n) \subset \mathrm{O}(kn)$  (where  $k = 1, 2, 4, 8$  depending on the field) and  $\partial\mathbf{H}^n$  is a sphere of real dimension  $(kn - 1)$ .  $\mathbb{Z}$

The following proposition is also known as the “cutting lemma” in a paper of Falbel and Gusevskii [FG94, th. 2.3], in which they directly refer to a paper of Kulkarni and Pinkall [KP86, th. 4.2]. A proof can be found in [KP86]. This result will be of great use in the proofs of Fried’s theorem III.1 and theorem IV.1.

**PROPOSITION II.7** *Let  $M$  be a closed  $(\mathrm{PU}(n, 1), \partial\mathbf{H}^n)$ -manifold. Denote  $\Gamma$  the holonomy group and  $\Omega = \partial\mathbf{H}^n - D(\tilde{M})$ . Suppose that  $L(\Gamma) \subset \Omega$  and that  $L(\Gamma)$  has at least two points. Then  $D$  is a covering map onto a connected component of  $\partial\mathbf{H}^n - L(\Gamma)$ .*

### III — FRIED’S THEOREM AND SIMILARITY STRUCTURES

The goal of this section is to show the following theorem, initially stated and shown by Fried for the real case  $\mathbf{F} = \mathbf{R}$  in [Fri80]. The complex case  $\mathbf{F} = \mathbf{C}$  was proved by Miner in [Min90]. Later, a different proof (but still in the initial ideas of Fried) of the real case was given by Matsumoto in his survey [Mat92]. An analytic proof of the real case was done by [BH99]. The quaternionic case  $\mathbf{F} = \mathbf{H}$  was addressed by Kamishima in [Kam99]. It seems that the octonionic case  $\mathbf{F} = \mathbf{O}$  has not been proved yet. We will simultaneously prove the theorem for all the fields  $\mathbf{F}$  considered.

**THEOREM III.1** *Let  $M$  be a closed  $(\mathrm{Sim}(\mathcal{N}), \mathcal{N})$ -manifold. If the developing map  $D: \tilde{M} \rightarrow \mathcal{N}$  is not a cover onto  $\mathcal{N}$ , then the holonomy subgroup  $\Gamma$  fixes a point in  $\mathcal{N}$  and  $D$  is in fact a covering onto the complement of this point.*

The different ideas of the proof come from [Fri80], [Min90] and [Mat92]. The ideas about convexity of Fried and Miner (also in Carrière’s work [Car89]) forged the necessity of the second section and the ideas for the first arguments of the theorem’s proof. Matsumoto’s ideas helped to find the last arguments.

#### III.1 A SHORTCUT: DISCRETE HOLONOMY AND AUTOSIMILARITY

A consequence of Fried’s theorem is that *for any similarity structure on a closed manifold, the holonomy is discrete*. The converse also holds.

**PROPOSITION III.2** *Suppose that  $\Gamma \subset \mathrm{Sim}(\mathcal{N})$  is a discrete subgroup. Then either  $L(\Gamma) = \emptyset$ , or  $L(\Gamma) = \{\infty\}$  or  $L(\Gamma) = \{\infty, a\}$  for some  $a \in \mathcal{N}$ .*

This can be compared with Matsumoto [Mat92, lemma 4.20].

**PROOF** Suppose that  $L(\Gamma)$  is neither  $\emptyset$  nor  $\{\infty\}$ . Since  $\Gamma \subset \mathrm{Sim}(\mathcal{N})$ , we have  $\infty \in L(\Gamma)$ . Indeed, if not, then  $\Gamma \subset M$  but  $L(M) = \emptyset$ .

Now, let  $f, g$  be such that  $f^n(x) \rightarrow a$  and  $g^n(x) \rightarrow b$ . This hypothesis can be made because if  $L(\Gamma) \neq \{\infty\}$  then there exist  $f, g$  with dilatation factors different from 1. If  $a = b$  for all choices  $f, g$  then  $\Gamma$  fixes  $a$  and it follows that  $L(\Gamma) = \{\infty\}$  or  $L(\Gamma) = \{a, \infty\}$ . So by absurd, suppose  $a \neq b$ .

Denote  $f(x) = \lambda P(x) + c$  and  $g(x) = \mu Q(x) + d$ . Take  $h = g \circ f \circ g^{-1}$ , we see that  $h$  fixes  $g(a)$ . Set also the sequence  $h_n = f^n \circ h \circ f^{-n}$ . The fixed point of  $h_n$  is  $f^n(g(a))$  tending to  $a$ .

$$\begin{aligned} g \circ f \circ g^{-1}(x) &= \mu Q(\lambda P(\mu^{-1} Q^{-1}(x) + d') + c) + d \\ &= \lambda Q P Q^{-1}(x) + e \end{aligned} \tag{28}$$

$$f^n \circ h \circ f^{-n}(x) = \lambda P^n Q P Q^{-1} P^{-n}(x) + c_n \tag{29}$$

Denote  $A_n = P^n Q P Q^{-1} P^{-n}$ . Since  $A_n \in M$  which is compact, we can extract a subsequence converging to  $A$ . Now, the constant  $c_n$  must tend to  $c$  since the fixed point tends to  $a$ . Hence,  $h_n(x)$  accumulates to  $\lambda A(x) + c$ , contradicting the discreteness of  $\Gamma$ .  $\mathbb{X}$

From this property it is not difficult to retrieve Fried's theorem fixed point property. If  $M$  is not complete, then  $L(\Gamma)$  is neither  $\emptyset$  nor  $\{\infty\}$ . Indeed, there must be an element of  $\Gamma$  with dilatation factor different from 1, giving a limit point in  $\mathcal{N}$  (otherwise the structure is complete, since there would exist an invariant riemannian metric). Therefore by discreteness,  $L(\Gamma)$  is  $\{a, \infty\}$  and  $a$  must be fixed. For closed manifolds, it is then possible to retrieve the full Fried's theorem with the fixed point property.

Of course, the hypothesis that  $\Gamma$  is discrete is highly non trivial, and this is why the proof of Fried's theorem is important. However, the author wishes to emphasize the fact that it remains true for any discrete subgroup (even if the manifold considered is not compact). In particular, it is possible to prove that any group with the property that *if a point of  $L(\Gamma)$  is totally fixed by  $\Gamma$  then  $L(\Gamma)$  is elementary (i.e. has at most two points)* enables to prove that  $L(\Gamma)$  is autosimilar when it is not elementary. For take  $p \in L(\Gamma)$  and  $U$  an open neighborhood of  $p$ . The complement  $\partial \mathbf{H}^n - \Gamma \cdot U$  is closed and invariant. Hence it must be at most a single point totally fixed by  $\Gamma$  since  $L(\Gamma)$  contains  $p$  and would be contained in this complement if there were more than one point. Now  $L(\Gamma)$  can in fact not have a totally fixed point in  $\partial \mathbf{H}^n - \Gamma \cdot U$  since we supposed that  $L(\Gamma)$  is infinite and verifies the property emphasized before. Therefore  $\Gamma \cdot U$  covers  $L(\Gamma)$ , and this shows the autosimilarity property.

### III.2 THE GEOMETRY OF $\mathcal{N}$

Some facts about the geometry of  $\mathcal{N}$  will be needed. In the real case,  $\mathcal{N}$  is the Euclidean space endowed with its similarities. The advantage is that the Euclidean space  $\mathbf{R}^n$  is flat and we are allowed to state  $\exp_x(v) = x + v$ .

It is still true in general. First,  $\mathcal{N}$  is a 2-nilpotent Lie group. It follows that the Lie algebra  $\mathfrak{n}$  of  $\mathcal{N}$  is 2-nilpotent and the exponential map is a diffeomorphism between  $\mathfrak{n}$  and  $\mathcal{N}$ . The real vector space  $\mathfrak{n}$  is to be thought as a global coordinate system of  $\mathcal{N}$ .

Geodesics of  $\mathcal{N}$  can be described in explicit terms. Let  $\omega$  be the Maurer-Cartan form of  $\mathcal{N}$ , i.e.  $\omega : T\mathcal{N} \rightarrow \mathfrak{n}$  is a 1-form constant on left-invariant vector fields. Geodesics  $\gamma : I \rightarrow \mathcal{N}$  are smooth curves such that  $\gamma^* \omega$  is constant. Geodesics from  $0 \in \mathcal{N}$  are given by  $\exp(tv)$  for any  $v \in \mathfrak{n}$ . If  $\exp(p) \in \mathcal{N}$  is any other point then the geodesics issued from  $p$  are given by left multiplication:  $\exp(p) \exp(tv)$ .

Note that in the coordinates of  $\mathfrak{n}$  given by the exponential, every geodesic is a straight line: from 0 it is clear since geodesics are given by  $\exp(tv)$  corresponding to



the coordinates  $tv$ . If  $\exp(p) \in \mathcal{N}$  then the Campbell-Hausdorff formula gives

$$\exp(p) \exp(tv) = \exp\left(p + tv + \frac{1}{2}[p, tv]\right) \quad (30)$$

hence coordinates of geodesics issued from  $p$  are given by  $p + tv + \frac{1}{2}[p, tv]$  which is again a straight line. This is a fact restricted to the 2-nilpotent groups, in general the Campbell-Hausdorff formula gives a polynomial which is not affine.

We will denote by  $\exp_{\exp(p)}(tv)$  or  $\exp_p(tv)$  the product  $\exp(p) \exp(tv)$  or  $p \exp(tv)$  depending whereas  $p \in \mathfrak{n}$  or  $p \in \mathcal{N}$ . Since geodesics are given by left translations, we can still write  $\exp_x(v) = x +_{\mathcal{N}} v$  as in the Euclidean case. We will be careful not to write  $\exp_x(tv) = x + tv$  since  $tv$  could indicate the dilatation by a factor  $t$  as in the group  $\mathcal{N}$  and not linearly as in the tangent space  $\mathfrak{n}$ .

Similarity transformations preserve the geodesic structure: if  $\rho \in \text{Sim}(\mathcal{N})$  and if  $\gamma$  is geodesic, then  $\rho\gamma$  is again geodesic. This can be checked manually since the group  $\text{Sim}(\mathcal{N})$  is well known.

The space  $\mathcal{N}$  is locally convex. The open balls that will be defined in the next section provide examples of arbitrary small open convex subsets.

If  $K$  is any manifold with a similarity structure, then we can pullback the geodesic structure on  $K$ . To be more specific, a curve  $\gamma : I \rightarrow K$  is a geodesic if and only if  $D \circ \gamma$  is a geodesic of  $\mathcal{N}$ . This construction implies the equivariance

$$D(\exp_x(v)) = \exp_{D(x)}(dD \cdot v) = D(x) + dD(v). \quad (31)$$

The exponential map at  $x$  in  $K$  is defined on an open subset. By definition, we will say that this is the *visible* open subset, denoted by  $V_x \subset T_x K$ . This open subset is non empty since  $D$  is a local diffeomorphism.

We are now interested in the case  $K = \tilde{M}$ . The following construction enables to get arguments about  $V_x$ . Let  $p \in \tilde{M}$  and  $g \in \pi_1(M)$ . Suppose that there exists  $v$  such that  $\exp_p(v) = g \cdot p$ . Since  $V_p$  is an open neighborhood of  $v$ , and since  $\exp$  is a local diffeomorphism, we can reduce  $V_p$  to a neighborhood  $W_p$  of  $v$  on which  $\exp$  is a diffeomorphism. Therefore, there exists an open neighborhood  $U_p$  of the origin in  $T_p \tilde{M}$  such that the application  $G$  defined on  $U_p$  by

$$G(u) = (\exp_p^{-1} \circ g \circ \exp_p)(u) \quad (32)$$

is well defined and verifies  $G(0) = v$  and  $G(U_p) = W_p$ . Remark that the maps  $g$  and  $G$  are equivariant:

$$\exp_p(G(u)) = g \cdot \exp_p(u). \quad (33)$$

We are now interested to extend  $G$  to the full tangent space  $T_p \tilde{M}$ . To do so, we need to treat the difficulty of the exponential of a non-visible vector. This is done by looking through the developing map.

As we did before, we can set  $\exp_x(w) = x + w$  for  $x \in \tilde{M}$  and  $w \in \mathfrak{n}$  by looking through the developing map ( $x \in \tilde{M}$  is sent to  $z \in \mathcal{N}$ , the tangent spaces are identified and any tangent space in  $\mathcal{N}$  is identified to  $\mathfrak{n}$  by the parallel transport). For convenience we will denote by  $p + V_p$  the image  $\exp_p(V_p)$  of the exponential of all visible vectors.

**LEMMA III.3** *For any  $w_1, w_2 \in T_p \tilde{M}$ , if  $D(p) + dD(w_1) = D(p) + dD(w_2)$  then  $w_1 = w_2$ . In particular  $D$  restricted to  $p + V_p$  is injective since  $D(p + v) = D(p) + dD(v)$ .*

**PROOF** If  $D(p) + dD(w_1) = D(p) + dD(w_2)$  then by unicity of the geodesics in  $\mathcal{N}$  we deduce  $dD(w_1) = dD(w_2)$ . Since  $D$  is a local diffeomorphism, this implies  $w_1 = w_2$ .  $\square$

In  $\mathcal{N}$  every exponential is well defined. By the preceding lemma, what happens in  $V_p$  is not different from what happens in the developing map. This is why we define

$$G(u) = dD_p^{-1} \circ \exp_{D(p)}^{-1} \circ \rho(g) \circ \exp_{D(p)} \circ dD_p(u). \quad (34)$$

With  $v$  as before, it remains true that  $G(0) = v$ . Again, for any  $w \in V_p \cap G^{-1}(V_p)$

$$\exp_p(G(w)) = g \cdot \exp_p(w). \quad (35)$$

For, recall that by the lemma the developing map is injective on  $p + V_p$ . We conclude by the following computation.

$$D(g \cdot \exp_p(w)) = \rho(g) \circ D(\exp_p(w)) \quad (36)$$

$$D(\exp_p(G(w))) = \exp_{D(p)}(dD \cdot G(w)) \quad (37)$$

$$= \rho(g) \circ \exp_{D(p)}(dD_p \cdot w) \quad (38)$$

$$= \rho(g) \circ D(\exp_p(w)) \quad (39)$$

The following proposition is crucial to the study. We will recall the notations.

**PROPOSITION III.4** *Let  $p \in \tilde{M}$  and  $v \in V_p$  such that there exists  $g \in \pi_1(M)$  verifying  $g \cdot p = p + v$ . The map  $G$  exponentially equivariant with  $g$  is defined by*

$$G(u) = dD_p^{-1} \circ \exp_{D(p)}^{-1} \circ \rho(g) \circ \exp_{D(p)} \circ dD_p(u). \quad (40)$$

*Suppose that  $U$  is a convex subset of  $V_p$  and contains  $v = G(0)$ . For any  $w \in T_p \tilde{M}$ , if  $G(w) \in U$  then  $w \in V_p$ .*

**PROOF** Suppose that  $U$  verifies the hypotheses. For  $t < \epsilon$  small enough  $tw$  is visible and  $G(tw)$  is close to  $v$  and is hence visible. Therefore  $tw \in V_p \cap G^{-1}(V_p)$ . We know by equation (35) that

$$\exp_p(G(tw)) = g \cdot \exp_p(tw). \quad (41)$$

Since  $\rho(g)$  transforms geodesics into geodesics, it transforms  $tw$  onto a geodesic from  $G(0)$  to  $G(w)$ .

Since  $G(0), G(w) \in U$  by assumption, the left member is well defined for any  $t \in [0, 1]$  because  $U$  is convex. The right member is defined for  $t$  in  $[0, \epsilon[$ . But since the left member is always defined  $[0, \epsilon[$  must be closed in  $[0, 1]$ , therefore equal to  $[0, 1]$ . It follows that  $w$  is visible.  $\mathbb{X}$

We will now give the first argument of Fried's theorem's proof. The hypothesis to keep in mind is that the similarity structure does not give a covering onto for  $D$ . The following ideas of convexity properties can be related to the work of Carrière in [Car89]. Some parts are exposed in [Min90].

Let  $C$  be an open subset of  $\tilde{M}$ . We will say that  $C$  is a *convex* subset if  $D|_C$  is a diffeomorphism with convex image in  $\mathcal{N}$ . Convexity in  $\mathcal{N}$  is the property of containing every geodesic segment.

**LEMMA III.5** *Let  $p \in \tilde{M}$  and  $0 \in C \subset V_p$  such that  $\exp_p(C)$  is convex. Let  $g \in \pi_1(M)$ . Then  $g \exp_p(C)$  is a convex subset containing  $gp$ .*

**PROOF** Since  $g$  transforms geodesics into geodesics, it sends a convex to a convex.  $\mathbb{X}$

Remark that by the following proposition, we furthermore have that  $g \exp_p(C) \subset gp + V_{gp}$ .

**PROPOSITION III.6** *We have the following properties.*

- (1) *If  $p \in C$  with  $C \subset \tilde{M}$  convex, then  $C \subset p + V_p$ .*
- (2) *If  $C_1, C_2$  are convex in  $\tilde{M}$  with a non empty intersection, then  $D|_{C_1 \cup C_2}$  is injective.*

(3) If  $\tilde{M}$  contains  $p$  such that  $p + V_p$  is convex, then  $p + V_p = \tilde{M}$ .

PROOF In order.

- (1) If  $z \in D(C)$  then there exists  $w \in \mathfrak{n}$  such that  $z = D(p) + w$ . Let  $v = (dD)^{-1}(w)$ . Then  $p + tv$  is well defined and belongs to  $C$  for  $t$  small enough. But  $D(p) + dD_p(tv)$  is well defined and in  $D(C)$  for all  $t \leq 1$ . Hence  $p + tv$  is well defined for  $t = 1$  by taking  $D^{-1}(D(p) + dD_p(tv))$ .
- (2) Take  $p \in C_1 \cap C_2$  and  $z = D(p)$ . If  $D(x_1) = D(x_2)$  for  $x_1 \in C_1$  and  $x_2 \in C_2$  then the geodesic segment from  $z$  to  $D(x_1)$  is the same that joins  $z$  to  $D(x_2)$ , hence for the same direction. Hence  $x_1 = x_2$ .
- (3) It suffices to show that  $p + V_p$  is closed in  $\tilde{M}$  since it is already open and non empty. Let  $y$  be in the adherence of  $p + V_p$ . Let  $C \subset y + V_y$  be a convex subset containing  $y$  (it can always be constructed since  $D$  is a local diffeomorphism). Then  $C \cap p + V_p$  has a non empty intersection, hence  $D$  is injective on  $C \cup p + V_p$ . In  $D(C \cup p + V_p)$  there exists  $v$  such that  $D(q) = D(p) + dD_p(v)$ . The geodesic  $D^{-1}(D(p) + dD_p(tv))$  is well defined and in  $p + V_p$  for  $t < 1$  by hypothesis. At  $t = 1$  the point is well defined by injectivity of  $D$ , is equal to  $q$  and belongs to the same geodesic, since for  $t$  large enough it belongs to  $C$ . Therefore  $q \in p + V_p$ .

✎

Remark that this last property shows that if  $p + V_p$  is convex, then  $D$  is a diffeomorphism.

**COROLLARY III.7** Suppose that  $D$  is not a covering map onto  $\mathcal{N}$  (hence not a diffeomorphism), then  $V_p$  is never equal to  $T_p\tilde{M}$ .

PROOF The tangent space  $T_p\tilde{M}$  is convex since  $dD_p$  is an isomorphism. If  $V_p = T_p\tilde{M}$ , then  $p + V_p = \tilde{M}$  and  $D(p + V_p) = D(p) + \mathfrak{n} = \mathcal{N}$ . It follows that  $D : \tilde{M} \rightarrow \mathcal{N}$  is a diffeomorphism, hence a covering.

✎

### III.3 PROOF OF FRIED'S THEOREM

We recall the hypotheses. The manifold  $M$  is a closed  $(\text{Sim}(\mathcal{N}), \mathcal{N})$ -manifold such that the developing map  $D : \tilde{M} \rightarrow \mathcal{N}$  is not a covering map.

Recall that we set a distance function  $d_{\mathcal{N}}(x, y) = \| -x + y \|$  from the pseudo-norm  $\| \cdot \|$  which is compatible with dilatations:  $\| \lambda x \| = |\lambda| \| x \|$ . The triangle inequality also remains true. This distance function is chosen so it is left-invariant:  $d_{\mathcal{N}}(a + x, a + y) = d_{\mathcal{N}}(x, y)$ . In particular, we can define the open ball of radius  $R$  centered in 0 to be

$$B(0, R) = \{x \in \mathcal{N} \mid d_{\mathcal{N}}(0, x) < R\}.$$

And in general, the open ball of radius  $R$  centered in  $p$  is given by the left translation of  $B(0, R)$  to  $p$ .

The preceding corollary shows that for every  $p \in \tilde{M}$ , the image  $D(p + V_p) =: D(\exp_p(V_p))$  is never equal to  $\mathcal{N}$ . Hence, for each  $p \in \tilde{M}$ , there exists an open subset  $B_p \subset V_p \subset T_p\tilde{M}$  such that the image  $D(p + B_p)$  is the maximal open ball in  $D(p + V_p)$  centered in  $D(p)$ .

We let

$$r : \tilde{M} \rightarrow ]0, +\infty[ \quad (42)$$

be the map that associates to  $p \in \tilde{M}$  the radius of the ball  $D(p + B_p)$  in  $\mathcal{N}$ .

**LEMMA III.8** For  $p \in \tilde{M}$  and  $q \in p + B_p$ ,

$$r(p) \leq r(q) + d_{\mathcal{N}}(D(p), D(q)) \quad (43)$$

and therefore  $r$  is a local contraction.

Furthermore, if  $g \in \pi_1(M)$  then  $r(gp) = \lambda(g)r(p)$  with  $\lambda(g)$  the dilatation factor of the holonomy transformation  $\rho(g) \in \text{Sim}(\mathcal{N})$ .

PROOF Let  $p \in \tilde{M}$ . and let  $q \in p + B_p$ . By proposition III.6,  $p + B_p \subset q + V_q$ . Let  $v \in \partial B_q$  such that  $q + v$  is not defined. Then  $D(q) + dD_q(v)$  does not belong to  $D(q + V_q)$  hence does not belong to  $D(p + B_p)$  which is precisely an open ball of radius  $r(p)$ . Therefore

$$r(p) \leq d_{\mathcal{N}}(D(p), D(q) + dD_q(v)) \quad (44)$$

$$\leq d_{\mathcal{N}}(D(q), D(q) + dD_q(v)) + d_{\mathcal{N}}(D(p), D(q)) \quad (45)$$

$$\leq r(q) + d_{\mathcal{N}}(D(p), D(q)). \quad (46)$$

To prove the second part, we prove that  $\rho(g)$  transforms  $D(p + B_p)$  into  $D(gp + B_{gp})$ . If that is true then for  $v \in \partial D(p + B_p)$ , we have  $\rho(g)v \in \partial D(gp + B_{gp})$  and therefore

$$r(gp) = d_{\mathcal{N}}(D(gp), \rho(g)v) = d_{\mathcal{N}}(\rho(g)D(p), \rho(g)v) \quad (47)$$

$$= d_{\mathcal{N}}(0, \rho(g)(-D(p) + v)) \quad (48)$$

$$= \lambda(g)d_{\mathcal{N}}(0, -D(p) + v) \quad (49)$$

$$= \lambda(g)d_{\mathcal{N}}(D(p), v) = \lambda(g)r(p). \quad (50)$$

In fact, it suffices to prove that  $\rho(g)D(p + B_p) \subset D(gp + B_{gp})$  since with  $g^{-1}$  we would get  $\rho(g)^{-1}D(gp + B_{gp}) = \rho(g^{-1})D(gp + B_{gp}) \subset D(p + B_p)$ , and by applying  $\rho(g)$  on both ends, we get  $D(gp + B_{gp}) \subset \rho(g)D(p + B_p)$ .

By lemma III.5,  $g$  sends  $p + B_p$  into a convex subset containing  $gp$ , and by proposition III.6 this convex is contained in  $gp + V_{gp}$ . But  $\rho(g)$  preserves open balls, hence  $\rho(g)D(p + B_p)$  is an open ball contained in  $D(gp + B_{gp})$  by maximality of  $B_{gp}$ .  $\mathbb{X}$

The equivariance between  $r$  and  $\lambda$  allows a sense of length in  $M$  which will be invariant by the holonomy group.

In  $\tilde{M}$  we set

$$d_{\tilde{M}}(p_1, p_2) = \frac{d_{\mathcal{N}}(D(p_1), D(p_2))}{r(p_1) + r(p_2)} \quad (51)$$

which is  $\pi_1(M)$ -invariant.

Let  $p \in \tilde{M}$  and let  $\epsilon > 0$ . We will describe open balls of radius  $\epsilon$  in  $\tilde{M}$  by locally looking at the pseudo-distance function  $d_{\tilde{M}}$  on couples in  $(p, p + B_p)$ . On this set,  $D$  is injective. Also, by lemma III.8, the function  $r$  is contracting. This gives

$$d_{\tilde{M}}(p, x) = \frac{d_{\mathcal{N}}(D(p), D(x))}{r(p) + r(x)} \quad (52)$$

$$\implies d_{\tilde{M}}(p, x) \geq \frac{d_{\mathcal{N}}(D(p), D(x))}{2r(x) + d_{\mathcal{N}}(D(p), D(x))} \quad (53)$$

hence if we suppose  $d_{\tilde{M}}(p, x) < \epsilon$  with  $\epsilon$  small enough, we get

$$\frac{d_{\mathcal{N}}(D(p), D(x))}{2r(x) + d_{\mathcal{N}}(D(p), D(x))} < \epsilon \quad (54)$$

$$\implies \frac{d_{\mathcal{N}}(D(p), D(x))}{r(x)} < \frac{2\epsilon}{1 - \epsilon}. \quad (55)$$

If  $r(p) \geq r(x)$  then the same inequality is true for  $r(p)$  instead of  $r(x)$ . If  $r(p) < r(x)$ , then  $p \in x + B_x$  ( $p$  is visible from  $x$  and lies inside the ball since it is closer to  $x$  than the boundary at distance  $r(x)$ ) and by repeting the same argument for  $(x, p)$  we get

the preceding inequality with  $r(p)$  instead of  $r(x)$ . In either cases

$$\frac{d_{\mathcal{N}}(D(p), D(x))}{r(p)} < \frac{2\epsilon}{1-\epsilon}. \quad (56)$$

Conversely by using  $r(x) \geq r(p) - d_{\mathcal{N}}(D(p), D(x))$

$$d_{\tilde{M}}(p, x) \leq \frac{d_{\mathcal{N}}(D(p), D(x))}{2r(p) - d_{\mathcal{N}}(D(p), D(x))} \quad (57)$$

hence for  $\epsilon > 0$

$$\frac{d_{\mathcal{N}}(D(p), D(x))}{2r(p) - d_{\mathcal{N}}(D(p), D(x))} < \epsilon \quad (58)$$

$$\implies \frac{d_{\mathcal{N}}(D(p), D(x))}{r(p)} < \frac{2\epsilon}{1+\epsilon} \quad (59)$$

This shows that for  $\epsilon$  small enough, the ball

$$B_{\tilde{M}}(p, \epsilon) := \{x \in p + B_p \mid d_{\tilde{M}}(p, x) < \epsilon\} \quad (60)$$

has an approximation in its developing image:

$$\left\{ \frac{d_{\mathcal{N}}(D(p), D(x))}{r(p)} < \frac{2\epsilon}{1+\epsilon} \right\} \subset D(B_{\tilde{M}}(p, \epsilon)) \subset \left\{ \frac{d_{\mathcal{N}}(D(p), D(x))}{r(p)} < \frac{2\epsilon}{1-\epsilon} \right\} \quad (61)$$

is contained in an open of  $\tilde{M}$ . Hence those open balls provide the same basis for the topology of  $\tilde{M}$ . This means that  $d_{\tilde{M}}(p, x) < \epsilon$  is true when in the ball  $D(p + B_p)$  normalized by the radius  $r(p)$ , the distance between  $x$  and  $p$  is less than  $\approx 2\epsilon$ .

If  $g \in \pi_1(M)$ , then by lemma III.5 and proposition III.6 (compare also with the proof of lemma III.8 where we proved that  $g(p + B_p) = gp + B_{gp}$ ),  $gB_{\tilde{M}}(p, \epsilon)$  is a subset of  $gp + B_{gp}$ . The distance function  $d_{\tilde{M}}$  being  $\pi_1$ -invariant, this shows that

$$\forall g \in \pi_1(M), gB_{\tilde{M}}(p, \epsilon) = B_{\tilde{M}}(gp, \epsilon). \quad (62)$$

In  $M$ , we can define a system of open neighborhoods by projecting the previously constructed balls of  $\tilde{M}$ .

$$B_M(x, \epsilon) := \pi(B_{\tilde{M}}(p, \epsilon)), p \in \pi^{-1}(x). \quad (63)$$

For  $\epsilon$  small enough, the ball  $B_M(x, \epsilon)$  is therefore a trivializing neighborhood of  $x$ , and this system of open balls gives the same topology for  $M$  as the original one.

We will now construct holonomy transformations which will be very contracting, with a common center point and with no rotation. The idea is to take  $v \in \partial B_p$  such that  $p + v$  is not defined and to compare with  $D(p) + dDv$  in  $\mathcal{N}$  where it must be defined. The holonomy transformations will be centered in  $D(p) + dDv =: z + w$ . See figure 1 for the global setting.

Consider  $p \in \tilde{M}$  such that  $\exp_p(tv)$  is defined for  $0 \leq t < 1$  but not for  $t = 1$ . The geodesic curve  $[c(t)] = \exp_p(tv)$  is an incomplete geodesic. In  $M$ , the corresponding curve  $c(t) = \pi([c(t)])$  is then an infinite long curve in a compact space. Hence, there is a recurrent point  $x \in M$ .

Let  $B_M(x, \epsilon)$  be a ball with  $\epsilon$  radius, for  $\epsilon > 0$  small enough such that  $B_M(x, \epsilon)$  is trivializing  $\pi: \tilde{M} \rightarrow M$ . Let  $0 < t_1 < \dots < t_n < \dots$  be entry times such that  $t_n \rightarrow 1$  and  $c(t_n) \in B_M(x, \epsilon)$  but  $c([t_n, t_{n+1}]) \not\subset B_M(x, \epsilon)$  (it just says that  $c$  exits  $B_M(x, \epsilon)$  before time  $t_{n+1}$ ). Since  $\epsilon$  is small enough, for each  $t_n$ , up to homotopy we can set uniquely  $\eta_n$  the segment from  $x$  to  $c(t_n)$  contained in  $B_M(x, \epsilon)$ .

By construction we have the following lemma.

**LEMMA III.9** *For any  $i$ ,  $[c(t_i)] \in p_i + B_{p_i}$  and  $d_{\tilde{M}}([c(t_i)], p_i) < \epsilon$ .*




Figure 1: The general setting.

**PROOF** By hypothesis and by the preceding discussion, since  $B_M(x, \epsilon)$  is trivializing, if  $c(t_i) \in B_M(x, \epsilon)$  then any lift of  $c(t_i)$  is in  $B_{\tilde{M}}(\hat{p}, \epsilon)$  for  $\hat{p} \in \pi^{-1}(x)$ . In particular  $[c(t_i)]$  is at distance at most  $\epsilon$  from  $p_i$  and lies in  $p_i + B_{p_i}$  by definition of  $B_{\tilde{M}}(p_i, \epsilon)$ .  $\boxtimes$

We set

$$g_{ij} = \eta_j^{-1} \star c(t)|_{[t_i, t_j]} \star \eta_i, \quad (64)$$

this is a family of transformations belonging to  $\pi_1(M, x)$ . We are now interested in  $g_{ij}$  acting on  $\tilde{M}$ . The path  $\tilde{g}_{ij}$  lifting  $g_{ij}$  sends  $p_i$  to  $p_j$  by construction. We denote by  $\gamma_{ij}$  the holonomy transformation  $\rho(g_{ij})$ , and we denote by  $\tilde{\gamma}_{ij}$  the image  $D(\tilde{g}_{ij})$  and by  $\tilde{\gamma}$  the image  $D([c(t)])$ .

The vector  $v$  initially chosen is sent to  $w$  by  $dD$  and  $z := D(p)$ . Each  $p_i$  is sent to  $z_i$  by  $D$ . See again figure 1.

**PROPOSITION III.10** *For  $i, j$  large enough with  $j \gg i$ , the transformation  $\gamma_{ij}$  is centered as close to  $z + w$  as desired, with a dilatation factor as close to 0 as desired and with an orthogonal part as close to identity as desired. (In other terms,  $\gamma_{ij}(x) = \lambda P(x - \beta) + \beta$  with  $\lambda \rightarrow 0$ ,  $P \rightarrow E$  and  $\beta \rightarrow z + w$ .)*

The idea of the proof consists in taking a closer look to figure 1. Suppose for a minute that every  $z_i$  is in fact  $\tilde{\gamma}(t_i)$ . Then the transformation  $\gamma_{ij}$  sends  $\tilde{\gamma}(t_i)$  to  $\tilde{\gamma}(t_j)$ . It is then clear that the transformation is very contracting and is centered in  $z + w$  with no rotation. It is the object of the proof to show that this approximation is correct. Note that in practice, a dilatation  $\lambda(x, y) = (\lambda x, \lambda^2 y)$  does not stabilize a geodesic.

**PROOF** Assume that  $\gamma_{ij}$  is an affine transformation given by

$$\gamma_{ij}(x) = \lambda P(x - \beta) + \beta, \quad (65)$$

with  $\lambda \in \mathbf{R}$ ,  $P \in U(n-1)$  and  $\beta \in \mathcal{N}$  constant. We start by showing that  $\beta \rightarrow z + w$  if  $P = E_n$  and if  $\lambda \rightarrow 0$ .

Denote  $(a_i, b_i) = \tilde{\gamma}(t_i)$  and  $(c, d) = \beta$ . By construction,  $\gamma_{ij}$  sends  $D(p_i) = z_i$  to  $D(p_j) = z_j$ . We will later show that  $z_i$  and  $\tilde{\gamma}(t_i)$  are very close for  $t_i$  large enough.

Hence, the additional and final hypothesis is that  $\gamma_{ij}$  sends  $(a_i, b_i)$  to  $(a_j, b_j)$ . It gives the following computation.

$$(a_j, b_j) = \lambda((a_i, b_i) - \beta) + \beta \quad (66)$$

$$= \lambda((a_i - c, b_i - d - \operatorname{Im}(a_i^* c)) + (c, d)) \quad (67)$$

$$= (\lambda(a_i - c) + c, \lambda^2(b_i - d - \operatorname{Im}(a_i^* c)) + d + \operatorname{Im}(\lambda(a_i - c) + c)^* c) \quad (68)$$

$$= ((1 - \lambda)c + \lambda a_i, (1 - \lambda^2)d + \lambda^2(b_i - \operatorname{Im}(a_i^* c)) + \operatorname{Im}(\lambda((a_i - c)^* c))) \quad (69)$$

It then suffices to resolve the values of  $c$  and  $d$ .

$$a_j = (1 - \lambda)(c) + \lambda a_i \quad (70)$$

$$c = (1 - \lambda)^{-1}(a_j - \lambda a_i), \quad (71)$$

$$b_j = (1 - \lambda^2)d + \lambda^2(b_i - \operatorname{Im}(a_i^* c)) + \operatorname{Im}(\lambda(a_i - c)^* c) \quad (72)$$

$$d = (1 - \lambda^2)^{-1}(b_j - \lambda^2(b_i - \operatorname{Im}(a_i^* c)) - \lambda \operatorname{Im}((a_i - c)^* c)) \quad (73)$$

We now use the fact that  $(a_i, b_i)$  is given by

$$\exp_z(t_i(a_0, b_0)) = z + (t_i a_0, t_i b_0) = (z_1 + t_i a_0, z_2 + t_i b_0 + t_i \operatorname{Im}(z_1^* b_0)), \quad (74)$$

where  $w = (a_0, b_0)$  and  $z = (z_1, z_2)$ . We now use the hypothesis  $\lambda \rightarrow 0$ , we get  $c \rightarrow a_j$  and  $b \rightarrow b_j$  which show that for  $t_i, t_j \rightarrow 1$  large enough, the point  $\beta = (c, d)$  is as close as desired to  $z + w$ .

It remains to show that we can indeed suppose that  $P = E_n$ ,  $\lambda \rightarrow 0$  and that  $z_i$  is arbitrary close to  $\tilde{\gamma}(t_i)$ .

Since  $U(\mathcal{N})$  is compact and since  $\gamma_{jk}\gamma_{ij} = \gamma_{ik}$ , the transformation  $P$  of  $\gamma_{ij}$  accumulates to the identity  $E_n$ .

To show  $\lambda \rightarrow 0$ , we will use lemma III.9. First we have by lemmas III.9 and III.8 and by definition of  $[c(t)]$

$$r(p_j) \leq r([c(t_j)]) + d_{\mathcal{N}}(z_j, \tilde{\gamma}(t_j)) \quad (75)$$

and by lemma III.9 and definition of  $d_{\tilde{M}}$

$$d_{\mathcal{N}}(z_j, \tilde{\gamma}(t_j)) < \epsilon (r(p_j) + r([c(t_j)])) \quad (76)$$

This gives

$$r(p_j) \leq (1 + \epsilon)r([c(t_j)]) + \epsilon r(p_j). \quad (77)$$

Remark  $r(p_j) = r(g_{ij}(p_i)) = \lambda(g_{ij})r(p_i)$ , it gives

$$\lambda(g_{ij})r(p_i) \leq (1 + \epsilon)r([c(t_j)]) + \epsilon \lambda(g_{ij})r(p_i) \quad (78)$$

$$\lambda(g_{ij}) \leq \frac{1 + \epsilon}{1 - \epsilon} \frac{r([c(t_j)])}{r(p_i)}. \quad (79)$$

Since the numerator tends to 0, for  $i$  fixed we get that  $\lambda \rightarrow 0$  for  $j$  tending to  $+\infty$ . So this is true for  $i, j$  large enough such that  $j \gg i$ .  $\mathbb{X}$

We now recall the construction made for proposition III.4. Let  $p \in \tilde{M}$ . If for  $g \in \pi_1(M)$ ,  $g \cdot p$  is visible from  $p$  by a vector in  $B_p$ , then if a vector  $u$  is such that  $G(u) \in B_p$ , then in fact  $p + u$  is visible from  $p$ , with  $G$  given by

$$G(u) = dD_p^{-1}(-z + \rho(g)(D(p) + dD_p(u))). \quad (80)$$

This allows to extend the set of vectors that can be seen from  $p$  further than  $B_p$ . By passing through the developing map (by lemma III.3), if  $dD_p G(u) \in dD_p \cdot B_p$  then it is true that  $G(u) \in B_p$ . Therefore if  $D(p + G(u)) \in D(p + B_p)$  then  $u$  is visible.

We now use the proposition. We first need to check that  $g_{ij}p$  is visible from  $p$ . We are only interested in the transformations  $g_{ij}$  for large  $i$  and  $j$ . We know that  $\gamma_{ij}z$  lies in the image of the ball  $D(p + B_p)$ , say with a vector  $w'$  such that  $\gamma_{ij}z = z + w'$ . Say  $v'$  corresponds to  $w'$  by  $(dD)^{-1}$ . It follows that  $g_{ij}p$  is visible from  $p$  by  $v' \in B_p$ .

The next proposition is inferred from this discussion.

**PROPOSITION III.11** *The exponential based in  $p$  is well defined on a “half-space” of  $T_p\tilde{M}$  given by*

$$H_p = \bigcup_{j \gg i \gg 0} dD^{-1}(\gamma_{ij}^{-1}(dD \cdot B_p)). \quad \mathbb{R} \quad (81)$$

Remark that in the Euclidean case  $\mathbf{F} = \mathbf{R}$  as in Fried’s proof, this half-space is given by  $\langle u, v \rangle < 1$ . We now take a closer look to the form of  $H_p$  in the general  $\mathcal{N}$  to prove that it is in fact a half-space.

For, we look at the images  $\gamma_{ij}^{-1}(dD \cdot B_p)$ . For more clarity, we will suppose that  $dD \cdot B_p$  is the unit ball centered in 0 in  $\mathcal{N}$  (after taking the exponential based in 0), denoted by  $B$ . In other words,  $(u, I) \in B$  if and only if  $\|u\|^2 + \|I\| < 1$ .

**LEMMA III.12** *The boundary  $\partial H_p \subset T_p\tilde{M}$  is through the developing map an affine subspace either vertical (meaning that  $(u, I) \in A \times \text{Im}(\mathbf{F})$  with  $A$  affine in  $\mathbf{F}^{n-2}$ ) or horizontal (meaning that  $(u, I) \in \mathbf{F}^{n-2} \times A$  with  $A$  affine in  $\mathbf{F}^{n-2}$ ).*

**PROOF** From what we know,  $g_{ij}$  is of the form

$$g_{ij}(x) = \lambda_{ij}(x - (z + w)) + (z + w) \quad (82)$$

with  $\lambda_{ij} \rightarrow 0$  and a small rotation that had been ignored. Therefore, we look at the transformations

$$g_\beta(\lambda)(x) = \lambda(x - \beta) + \beta \quad (83)$$

with  $\beta \in \partial B$  the center and  $\lambda > 0$  the dilatation factor.

The image set for  $x$  fixed is

$$\{\lambda(x - \beta) + \beta \mid \lambda > 0\} \quad (84)$$

and this is half a parabola (sometimes degenerated into a straight line). We will examine the intersection of this parabola with  $B$  for small  $\lambda$ . If this intersection is non empty then  $x$  will be visible from  $p$ . Taking coordinates  $x = (x_1, x_2)$  and  $\beta = (\beta_1, \beta_2)$ ,

$$g_\beta(\lambda)(x) = (\lambda(x_1 - \beta_1), \lambda^2(x_2 - \beta_2 + \text{Im}(x_1^* \beta_1))) + (\beta_1, \beta_2) \quad (85)$$

$$= (\lambda(x_1 - \beta_1) + \beta_1, \lambda^2(x_2 - \beta_2) + \beta_2 + \lambda(1 + \lambda)\text{Im}(x_1^* \beta_1)). \quad (86)$$

First, we examine the case  $\beta_1 \neq 0$ . Up to applying a rotation of the subgroup  $M$ , we can suppose that  $\beta_1 = (c, 0, \dots, 0)$  with  $c > 0$  real. Denoting  $x_1 = (x_1^1, x_1^2, \dots, x_1^{n-1})$ , we get

$$g_\beta(\lambda)(x) = (\lambda(x_1 - \beta_1) + \beta_1, \beta_2 - \lambda c \text{Im}(x_1^1) + \lambda^2(x_2 - \beta_2 - c \text{Im}(x_1^1))) \quad (87)$$

$$= (\lambda(x_1 - \beta_1) + \beta_1, \beta_2 - \lambda c \text{Im}(x_1^1) + o(\lambda)) \quad (88)$$

When  $\lambda \rightarrow 0$ , the condition  $g_\beta(\lambda)(x) \in B$  does not depend on the coordinate  $x_2$ . Hence  $\partial H_p$  is vertical (the condition is affine since it only involves  $\text{Im}(x_1^1)$ ).

Now we suppose  $\beta_1 = 0$ . Hence  $\beta_2 = \xi$  is unitary. The coordinates of the parabola then are

$$g_\beta(\lambda)(x) = (\lambda x_1, \lambda^2(x_2 - \xi) + \xi). \quad (89)$$



The norm of the first coordinate tends to 0 and the norm of the second is smaller than 1 if  $\operatorname{Re}((x_2 - \xi)^* \xi) < 0$ , i.e.  $\operatorname{Re}(x_2^* \xi) < 1$ . Therefore  $\partial H_p$  is horizontal since this condition does not depend on  $x_1$  and is affine in  $x_2$ .  $\mathbb{Z}$

**COROLLARY III.13** *For each  $p \in \tilde{M}$ ,  $H_p$  is convex.*

**PROOF** Indeed,  $dD_p \partial H_p$  is affine and of real codimension 1, hence it separates  $n$  in two connected components, each convex. One of them is  $dD_p H_p$ .  $\mathbb{Z}$

We now divide  $\partial H_p$  onto  $W_p \sqcup I_p$ , where  $W_p$  denotes the visible vectors and  $I_p$  the invisible vectors of  $\partial H_p$ .

**LEMMA III.14** *The image  $D(p) + dD_p(I_p)$  is locally constant (hence constant) following  $p$ , this image is denoted  $I$ . Furthermore,  $I$  is affine.*

**PROOF** If  $W_p = \emptyset$ , then  $I_p = \partial H_p$  is affine and its image by the developing map must be constant since  $p + V_p = \tilde{M}$  is convex (because  $H_p$  is convex and equal to  $V_p$  if  $\partial H_p$  is only constituted of invisible vectors, see proposition III.6).

Let  $v_p \in I_p$ . Suppose that  $u \in W_p$ . Let  $q = p + u$ . The point  $q$  has a half-space  $H_q$ . We show that  $D(p) + dD_p(v_p)$  belongs to  $D(q) + dD_q(\partial H_q)$ . This shows that  $D(p) + dD_p(I_q)$  is contained in the intersection of  $D(p) + dD_p(\partial H_p)$  with  $D(q) + dD_q(\partial H_q)$ . Since  $D(q) + dD_q(\partial H_q) \subsetneq D(p) + dD_p(\partial H_p)$ , such an intersection decreases the topological dimension. By repeating the argument for a new  $u$ , the image of  $I_p$  becomes constant following  $p$  and is affine.

Since  $p + H_p$  and  $q + H_q$  are convex and with non empty intersection, it follows that  $D$  is injective on  $(p + H_p) \cup (q + H_q)$ . Suppose that  $D(p) + dD_p(v_p)$  lies in  $D(q + H_q)$ , then this is locally true. Therefore,  $D(c(t)) = D(p + tv_p)$  is contained in  $D(q + H_q)$  for  $t \in ]T, 1[$ . By injectivity of  $D$ , this shows that  $c(t) = p + tv_p$  is contained in  $q + H_q$  for  $t \in ]T, 1[$  and this geodesic is defined for  $t = 1$  by hypothesis. But this contradicts that  $c(1)$  is not defined. This shows that  $D(p) + dD_p(v_p)$  does not belong to  $D(q + H_q)$ .

Take  $p'$  in  $p + H_p$  such that  $D(p) + dD_p(v_p)$  belongs to  $\partial D(p' + B_{p'})$ . Then there exists  $\rho(g_{ij})$  centered in  $D(p) + dD_p(v_p)$  very contracting with almost no rotation for  $i, j$  large enough.

If  $D(p) + dD_p(v_p)$  does not belong to  $D(q) + dD_q(\partial H_q)$ , then this last set has no fixed point under  $\rho(g_{ij})$ . Therefore,  $D(g_{ij}(q + H_q))$  is convex and contains  $D(q) + dD_q(\overline{B_q})$ . But  $g_{ij}(q + H_q)$  and  $q + H_q$  intersect, hence  $D$  is injective on the union. If  $c(t) \in q + B_q$  is a geodesic such that  $c(0) = q$  and  $c(1)$  is not defined, then this shows that  $c(t)$  is well defined in  $t = 1$ , since it is in  $g_{ij}(q + H_q)$ , absurd.  $\mathbb{Z}$

Now the end of Fried's theorem's proof consists in showing that the holonomy group  $\Gamma$  is discrete (compare with III.2 and with [Mat92]).

Since  $I$  is constant, it follows that  $D(\tilde{M})$  does not intersect  $I$  (since the exponential of a point is always locally defined and  $I$  is the boundary of every  $H_p$ ). Since  $I$  is non empty by construction, by the proposition II.7, it follows that the developing map is a covering onto its image, which is  $\mathcal{N} - I$ .

The main argument is the following lemma, which proof can already be found in the proof of II.7.

**LEMMA III.15** *If there exist  $f, g \in \operatorname{Sim}(N)$  such that  $\lambda(f) \neq 1$  and the fixed point of  $f$  is not fixed by  $g$ , then  $\langle f, g \rangle$  is not a discrete subgroup of  $\operatorname{Sim}(\mathcal{N})$ .*

Such applications are given by the various  $g_{ij}$  by changing the base point  $p$ . If  $I$  is not a single point, then two such maps  $f, g$  exist. We will show that it does not occur. If  $\mathcal{N} - I$  is simply connected, then the holonomy group must be discrete, which contradicts the preceding lemma.

So we suppose, that  $\mathcal{N} - I$  is not simply connected. This case occurs if  $I$  is an affine subspace of real codimension 2. Let  $H$  be an affine half-space with  $\partial H = I$ . By turning  $H$  around  $I$ , we find that the universal cover of  $\mathcal{N} - I$  is  $H \times \mathbf{R}$ .

Since  $I$  is invariant by the holonomy group  $\Gamma$ , and since  $\Gamma$  contains contractions (such as  $g_{ij}$ ) it follows that  $I$  is stable by dilatations. The subgroup of  $\text{Sim}(\mathcal{N})$  stabilizing  $I$  contains dilatations  $f, g$  with different fixed points. The lifted group  $\text{Sim}(I) \times \mathbf{R}$  contains  $\tilde{f}, \tilde{g}$  (essentially  $\tilde{f} \simeq f \times \{0\}$  since  $f = g_{ij}$  does not rotate much around  $I$ ) again contracting and with different fixed points.

We now have a lifting of  $(G, X)$ -structures

$$(\text{Sim}(I) \times \mathbf{R}, H \times \mathbf{R}) \rightarrow (\text{Sim}(I), \mathcal{N} - I) \quad (90)$$

and  $M$  gets a lifted  $(\text{Sim}(I) \times \mathbf{R}, H \times \mathbf{R})$ -structure. The developing map is given by the choice of a point: take  $p \in \tilde{M}$  and  $D(p) \in \mathcal{N} - I$ , we have to choose  $q \in H \times \mathbf{R}$  such that  $q$  is sent to  $D(p)$  by the covering  $H \times \mathbf{R} \rightarrow \mathcal{N} - I$ , denote  $q$  by  $D'(p)$ . The new developing map  $D'$  is now fully prescribed by  $D$  and  $D'(p)$ . Again,  $D'$  is a covering map since  $D$  is a covering map. For, take  $q(t)$  a path in  $H \times \mathbf{R}$ , based in  $q(0) = D'(p)$ . It is sent to a path in  $\mathcal{N} - I$  which is covered by a path  $\tilde{q}(t)$  in  $\tilde{M}$  since  $D$  is a covering map, and  $D'(\tilde{q}(t)) = q(t)$  by local triviality.

Since  $H \times \mathbf{R}$  is simply connected, the new holonomy group  $\Gamma'$  must be discrete. But this is again contradicted by the preceding lemma (slightly adapted).

This concludes Fried's theorem's proof since  $I$  must be constituted of a single point (which must be totally fixed by  $\Gamma$ ). Also by the proposition II.7 the rest of the theorem is shown.  $\square$

#### IV — COMPLETE STRUCTURES ON CLOSED MANIFOLDS

The aim of this final section is to show the following theorem. This is more classical and not too difficult once Fried's theorem is given. See [KP86].

**THEOREM IV.1** *Let  $M$  be a closed  $(\text{PU}(n, 1), \partial\mathbf{H}^n)$ -manifold. If  $D$  is not surjective then it is a covering onto its image. Furthermore,  $D$  is a covering on its image if, and only if,  $D(\tilde{M})$  is equal to a connected component of  $\partial\mathbf{H}^n - L(\Gamma)$ .*

The proof of this theorem is cut into two parts.

**PROPOSITION IV.2** *Let  $M$  be a closed  $(\text{PU}(n, 1), \partial\mathbf{H}^n)$ -manifold. If the developing map is not surjective, then it is a covering map on its image.*

**PROOF** Denote  $\Omega = \partial\mathbf{H}^n - D(\tilde{M})$ .

- If  $\Omega = \{a\}$ , then up to conjugation, we can suppose that  $\Omega = \{\infty\}$  and  $D(\tilde{M}) \subset \mathcal{N}$ . Since  $\Omega$  is fixed by the holonomy group, we get a similarity structure  $(\text{Sim}(\mathcal{N}), \mathcal{N})$  on  $M$ . Since  $M$  is closed, it verifies the hypotheses of Fried's theorem and hence must be complete. Otherwise, another point would be missed by the developing map.
- Suppose  $\{a\} \subsetneq \Omega$ . Then by minimality  $L(\Gamma) \subset \Omega$ . If  $L(\Gamma) = \emptyset$  then by lemma II.5, the developing map is a covering onto the full  $\partial\mathbf{H}^n$ , absurd. Hence  $L(\Gamma)$  consists of one or at least two points. If  $L(\Gamma) = \{a\}$  then we can suppose it is  $\infty$  and  $D(\tilde{M}) \subset \mathcal{N}$ . Again, by Fried's theorem and since  $L(\Gamma) = \{\infty\}$ , the developing map is in fact a covering onto  $\mathcal{N}$ , absurd. Hence,  $L(\Gamma)$  consists of at least two points and is contained in  $\Omega$ . The conclusion follows from proposition II.7.

$\square$

**PROPOSITION IV.3** *Let  $M$  be a closed  $(\mathrm{PU}(n, 1), \partial\mathbf{H}^n)$ -manifold. The developing map is a covering on its image if, and only if,  $D(\tilde{M})$  is equal to a connected component of  $\partial\mathbf{H}^n - L(\Gamma)$ .*

**PROOF** We start by supposing that  $D(\tilde{M})$  is equal to the full connected component of  $\partial\mathbf{H}^n - L(\Gamma)$  on which  $D(\tilde{M})$  is defined.

- If  $L(\Gamma) = \emptyset$  then by lemma II.5 the conclusion follows.
- If  $L(\Gamma) = \{a\}$  then  $M$  gets a similarity structure which must be complete by Fried's theorem III.1.
- If  $L(\Gamma)$  is constituted of at least two points, then by hypothesis the developing map avoids  $L(\Gamma)$  and the proposition II.7 concludes.

Now we suppose that the developing map is a covering onto its image.

- If  $D$  is a covering onto  $\partial\mathbf{H}^n$  or  $\partial\mathbf{H}^n - \{a\}$ , then  $D$  is a diffeomorphism and  $\Gamma$  is discrete. It must avoid  $L(\Gamma)$ . If  $D$  is a covering onto  $\partial\mathbf{H}^n$  then  $L(\Gamma) = \emptyset$ . If  $D$  is a covering onto  $\partial\mathbf{H}^n - \{a\}$  then  $L(\Gamma) \neq \emptyset$  by lemma II.5, hence  $L(\Gamma)$  is exactly  $a$ .
- Now suppose that  $D$  is a covering and avoids at least two points. Let  $\Omega = \partial\mathbf{H}^n - D(\tilde{M})$ . Then by minimality  $L(\Gamma) \subset \Omega$  and  $L(\Gamma)$  can not be  $\emptyset$  nor  $\{a\}$ , otherwise it would be a complete boundary manifold (by lemma II.5) or a complete similarity manifold (by Fried's theorem III.1) and  $\Omega$  would be equal to  $\emptyset$  or  $a$ . So  $L(\Gamma)$  contains at least two points and we conclude by proposition II.7.

⌘

## REFERENCES

- [All99] Daniel Allcock. “Reflection groups on the octave hyperbolic plane”. In: *Journal of Algebra* 213.2 (1999), pp. 467–498 (cit. on p. 5).
- [Bae02] John C. Baez. “The octonions”. In: *American Mathematical Society. Bulletin. New Series* 39.2 (2002), pp. 145–205 (cit. on p. 5).
- [BH99] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature*. Vol. 319. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1999 (cit. on pp. 6, 7).
- [Car89] Yves Carrière. “Autour de la conjecture de L. Markus sur les variétés affines”. In: *Inventiones Mathematicae* 95.3 (1989), pp. 615–628 (cit. on pp. 7, 10).
- [CG74] Shyan S. Chen and Leon Greenberg. “Hyperbolic spaces”. In: *Contributions to analysis (a collection of papers dedicated to Lipman Bers)* (1974), pp. 49–87 (cit. on pp. 5, 6).
- [Cow+91] Michael Cowling et al. “ $H$ -type groups and Iwasawa decompositions”. In: *Advances in Mathematics* 87.1 (1991), pp. 1–41 (cit. on p. 5).
- [Fal08] Elisha Falbel. “A spherical CR structure on the complement of the figure eight knot with discrete holonomy”. In: *Journal of Differential Geometry* 79.1 (2008), pp. 69–110 (cit. on p. 6).
- [FG94] Elisha Falbel and Nikolay Gusevskii. “Spherical CR-manifolds of dimension 3”. In: *Boletim da Sociedade Brasileira de Matemática. Nova Série* 25.1 (1994), pp. 31–56 (cit. on p. 7).
- [Fri80] David Fried. “Closed similarity manifolds”. In: *Commentarii Mathematici Helvetici* 55.4 (1980), pp. 576–582 (cit. on pp. 1, 7).

- [Har90] F. Reese Harvey. *Spinors and calibrations*. Vol. 9. Perspectives in Mathematics. Academic Press, Inc., Boston, MA, 1990 (cit. on p. 5).
- [Kam99] Yoshinobu Kamishima. “Geometric superrigidity of spherical pseudo-quaternionic manifolds”. In: *The Proceedings of the Conference on Geometric Structures on Manifolds (Seoul, 1997)*. Vol. 46. Lecture Notes Ser. Seoul Nat. Univ., Seoul, 1999, pp. 111–149 (cit. on pp. 1, 7).
- [KP03] Inkang Kim and John R. Parker. “Geometry of quaternionic hyperbolic manifolds”. In: *Mathematical Proceedings of the Cambridge Philosophical Society* 135.2 (2003), pp. 291–320 (cit. on p. 3).
- [KP86] Ravi S. Kulkarni and Ulrich Pinkall. “Uniformization of geometric structures with applications to conformal geometry”. In: *Differential geometry, Peñíscola 1985*. Vol. 1209. Lecture Notes in Math. Springer, Berlin, 1986, pp. 190–209 (cit. on pp. 7, 18).
- [Mat92] Shigenori Matsumoto. “Foundations of flat conformal structure”. In: *Aspects of low-dimensional manifolds*. Vol. 20. Adv. Stud. Pure Math. Kinokuniya, Tokyo, 1992, pp. 167–261 (cit. on pp. 7, 17).
- [Min90] Robert R. Miner. “Spherical CR manifolds with amenable holonomy”. In: *International Journal of Mathematics* 1.4 (1990), pp. 479–501 (cit. on pp. 1, 7, 10).
- [Rie82] Carl R. Riehm. “The automorphism group of a composition of quadratic forms”. In: *Transactions of the American Mathematical Society* 269.2 (1982), pp. 403–414 (cit. on p. 5).
- [Rie84] Carl R. Riehm. “Explicit spin representations and Lie algebras of Heisenberg type”. In: *Journal of the London Mathematical Society. Second Series* 29.1 (1984), pp. 49–62 (cit. on p. 5).
- [Thu97] William P. Thurston. *Three-dimensional geometry and topology. Vol. 1*. Ed. by Silvio Levy. Princeton Mathematical Series. Princeton University Press, Princeton, NJ, 1997 (cit. on pp. 6, 7).